





Please address all correspondence to the Director General Fa'amolemole fa'atuatusi uma feso'otaiga i le Fa'atonusili Aoao

Syndromic Surveillance Report

Epi-week 38: 16/09/24 - 22/09/24

Date of report: 23rd September, 2024 **Prepared by:** National Health Surveillance & IHR Division

Email: surveillance@health.gov.ws Tel: 66506/66507


1. Dengue like illness (DLI)

Case Definition: Fever (≥38°C) for at least 2 days, PLUS at least 2 of the following: Nausea or vomiting; Muscle or joint pain; Severe headache or pain behind the eyes; Rash; bleeding

Alert threshold: twice the average number of cases seen in the previous two weeks

The state of the s		
Epi-weeks	Epi-week 37: 09/09/24 - 15/09/24	Epi-week 38: 16/09/24 - 22/09/24
Total cases reported	2	6
Moving Average	6	1
Threshold	12	2

Figure 1. Number of DLI cases reported from all health facilities



Dengue like illness case count is below the alert threshold. The Ministry of Health highly recommends the public to continue the practice of preventative measures such as: clean-up and remove stagnant water sources around the house that are potential mosquito breeding sites; wear appropriate clothing to minimize mosquito bites; and use mosquito nets and repellents such as lotion and sprays. If you are affected, watch out for severe symptoms and seek medical attention immediately. Also monitor other people in the household for similar symptoms and present to the hospital for testing and health advice.










2. Influenza like illness (ILI)

Case Definition: An acute respiratory infection with a history of fever or measured fever of ≥ 38 C° and cough, with onset within the last 10 days

Alert threshold: Exceeding double the average in the previous 2 weeks

Epi-weeks	Epi-week 37: 09/09/24 - 15/09/24	Epi-week 38: 16/09/24 - 22/09/24
Total cases reported	1118	1189
Moving Average	1358	1283
Threshold	2715	2565

Figure 2. Number of ILI cases reported from all Health Facilities



The influenza like illness case count is below the alert threshold. However, we continue to observe an increase of people presenting to the hospital with flu-like symptoms, and the numbers remain high. The Ministry of Health still highly recommends the general public to continue the practice of preventative measures such as – wearing of masks especially when you're having flu-like symptoms such as cough and runny nose, hand hygiene and avoid large gatherings without masks. If you are feeling ill with flu-like symptoms, please see a nurse or doctor, stay home and take medications as prescribed.

3. Severe acute respiratory infection (SARI)

Case Definition: An acute respiratory infection with a history of fever or measured fever of ≥ 38 °C and cough, with onset within the last 10 days, AND requiring hospitalization

Alert threshold: 2 linked cases

Epi-weeks	Epi-week 37: 09/09/24 - 15/09/24	Epi-week 38: 16/09/24 - 22/09/24
Total cases reported	13	7
Moving Average	22.5	17
Threshold	2 linked cases	2 linked cases









Figure 3. Number of SARI cases reported from all health facilities




The severe acute respiratory infection cases reported have been investigated, and no known links were identified. Therefore, it does not meet the definition exceeding the threshold. Despite the threshold, the case count remains high. Therefore, the public is strongly advised to continue to practice preventative measures as mentioned above such as wearing of masks, hand hygiene and avoid crowded places.

4. Diarrhea

Case Definition: Three (3) or more loose or watery or bloody stools in 24hrs Alert threshold: more than twice the average of the last two weeks count				
Epi-weeks	Epi-week 37: 09/09/24 - 15/09/24	Epi-week 38: 16/09/24 - 22/09/24		
Total cases reported	151	115		
Moving Average	110.5	121.5		
Threshold	221	243		

Figure 4. Number of Diarrhea cases reported from all health facilities



Diarrhea case count remains below the alert threshold. The Ministry of Health advises the public to continue good hand hygiene or hand washing practices, boil water before drinking and ensure food preparation is safe and sanitary.

